

October 2004

Zero-curvature solutions of the one-dimensional Schrödinger equation

M. Belloni*

Physics Department

Davidson College

Davidson, NC 28035 USA

M. A. Doncheski†

Department of Physics

The Pennsylvania State University

Mont Alto, PA 17237 USA

R. W. Robinett‡

Department of Physics

The Pennsylvania State University

University Park, PA 16802 USA

(Dated: February 1, 2008)

Abstract

We discuss special $k = \sqrt{2m(E - V(x))/\hbar^2} = 0$ (*i. e.* zero-curvature) solutions of the one-dimensional Schrödinger equation in several model systems which have been used as idealized versions of various quantum well structures. We consider infinite well plus Dirac delta function cases (where $E = V(x) = 0$) and piecewise-constant potentials, such as asymmetric infinite wells (where $E = V(x) = V_0 > 0$). We also construct supersymmetric partner potentials for several of the zero-energy solutions in these cases. One application of zero-curvature solutions in the infinite well plus δ -function case is the construction of ‘designer’ wavefunctions, namely zero-energy wavefunctions of essentially arbitrary shape, obtained through the proper placement and choice of strength of the δ -functions.

PACS numbers: 03.65.Ge, 03.65.Ca, 03.65.Db

*Electronic address: mabelloni@davidson.edu

†Electronic address: mad10@psu.edu

‡Electronic address: rick@phys.psu.edu

I. INTRODUCTION

Many important quantum-mechanical systems, such as the hydrogen atom and the rotational or vibrational states of molecules, whose study contributed to the early success of quantum theory, are nicely coincident with a few of the handful of simple quantum-mechanical problems which can be solved easily, namely the Coulomb potential, angular momentum states, and the harmonic oscillator. While textbooks often treat other similarly tractable bound state problems, many of these models have only recently found experimental realizations. For example, the quantum mechanical problem of a particle ‘bouncing’ from a flat surface under the influence of a constant force (*e. g.* gravity) is a staple in many texts and is easily soluble in terms of Airy functions, but experimental evidence for the lowest stationary state of neutrons in Earth’s gravitational field has only recently been presented [1]. The creation of a wide variety of non-classical motional states (*e. g.* coherent and squeezed states) of ions in artificially produced harmonic traps [2] is another example of the experimental realization of specific ‘designer’ quantum wavefunctions. Asymmetric quantum wells (modeled by piecewise-constant potentials with one side at a higher constant potential, V_0 , than the other) are simple examples of systems which can exhibit both quasi-classical and decidedly wave-like behavior (for $E > V_0$, where the solutions are oscillatory in both parts of the well) as well as providing an easy introduction to quantum tunneling (when $E < V_0$, so that the wavefunctions on each side have opposite curvature). Besides being of pedagogical interest, experiments exciting coherent charge oscillations from just such asymmetric multiple quantum well structures have been reported [3], making use of both types of solutions in the produced wave packet.

Other idealized potential systems, such as the finite square well [4] and infinite square well with additional Dirac delta function (δ -function) potentials, either with Dirichlet [5] or periodic boundary conditions [6] - [9] have been also discussed, often in the context of wave packet revival behavior or bouncing/trapping. It has been suggested [8] that such systems can be experimentally realized using microwave analogs [10] or as the limit of highly localized quantum wells. In a number of such recent examples then, the emphasis has shifted from searching for physical systems with correspondingly tractable mathematical analogs, to producing experimentally realizable versions of model potentials which can then exhibit desired characteristics, including generating very specific quantum wavefunctions.

In this note, we focus on the observation that a number of the model systems mentioned above, besides exhibiting negative-curvature (oscillatory) and positive-curvature (exponential) energy eigensolutions, can also support zero-curvature (hereafter ZC) eigenstates, *i. e.* straight-line solutions. Such solutions are seldom, if ever, discussed in the literature, or considered in more general analyses of these idealized systems. Such ZC solutions of the 1D Schrödinger equation, for which $k = \sqrt{2m(E - V(x))/\hbar^2} = 0$ over a finite spatial interval, can be supported in asymmetric quantum wells consisting of piecewise constant potentials (where $E = V(x) = V_0 > 0$ in part of the well) or in otherwise free systems (where $E = V(x) = 0$) where additional δ -function interactions are present to accommodate the boundary conditions. We will consider the latter case below in some detail and briefly discuss aspects of the former case at the end. We will also make note of the fact that zero-energy solutions of the infinite square well plus δ -function case can be used to generate interesting supersymmetric partner potentials.

II. INFINITE WELL PLUS δ -FUNCTION INTERACTIONS

For the standard infinite square well (ISW) defined over the region $[0, a]$, zero-energy, and hence zero-curvature, solutions of the Schrödinger equation of the form $\psi(x) = A + Bx$ are formally possible. However the imposition of the boundary conditions $\psi(0) = 0 = \psi(a)$ yields the uninteresting result that $A = B = 0$, so that $\psi(x)$ vanishes identically. On the other hand, a waveform given by

$$\psi(x) = \begin{cases} 0 & \text{for } x \leq 0 \text{ and } a \leq x \\ Ax/c & \text{for } 0 \leq x \leq c \\ A(a-x)/(a-c) & \text{for } c \leq x \leq a \end{cases} \quad (1)$$

is a ZC solution of the Schrödinger equation which satisfies the boundary conditions at $x = 0$ and $x = a$ (and which is appropriately normalized if $A = \sqrt{3/a}$), but which has a discontinuity in $\psi'(x)$, a cusp, at $x = c$. This form can be accommodated by the addition of an appropriate strength (attractive) δ -function potential of the form $V_\delta(x; c) \equiv g\delta(x - c)$ since such a potential induces a discontinuity in the derivative given by

$$\psi'(c^+) - \psi'(c^-) = \left(\frac{2mg}{\hbar^2} \right) \psi(c). \quad (2)$$

For the waveform in Eqn. (1), the appropriate strength, g_0 , required to yield a ZC solution [11] is given by

$$\left(-\frac{A}{(a-c)}\right) - \left(\frac{A}{c}\right) = \left(\frac{2mg_0}{\hbar^2}\right) A \quad \text{or} \quad g_0 = -\frac{\hbar^2}{2m} \frac{a}{c(a-c)} \quad (3)$$

which is independent of the normalization, A , and only depends on the shape of the triangular waveform. The interaction strength required diverges as $c \rightarrow 0$ and $c \rightarrow a$ as expected since the resulting waveform has an increasingly rapid spatial variation, implying arbitrarily large kinetic energy, as the waveform is forced to jump almost discontinuously from zero to A (or from A to zero) near one of the two boundaries, as seen in Fig. 1.

This observation can be confirmed more quantitatively by calculating the expectation values of the potential and kinetic energies in this modified ISW system. The expectation value of the potential energy is given by

$$\langle V_\delta(x; c) \rangle = \int_0^a V_\delta(x; c) |\psi(x)|^2 dx = g_0 |\psi(c)|^2 = -\frac{3\hbar^2}{2mc(a-c)}. \quad (4)$$

The expectation value of kinetic energy can be expressed in one of two equivalent forms, namely

$$\langle \hat{T} \rangle = \int \psi^*(x) \left(\frac{\hat{p}^2}{2m} \right) \psi(x) dx = -\frac{\hbar^2}{2m} \int \psi^*(x) \frac{d^2\psi(x)}{dx^2} dx \quad (5)$$

or

$$\langle \hat{T} \rangle = \frac{\hbar^2}{2m} \int \left| \frac{d\psi(x)}{dx} \right|^2 dx \quad (6)$$

where the expression in Eqn. (6) is obtained from Eqn. (5) by using an integration by parts. Use of the form in Eqn. (6) immediately gives

$$\langle \hat{T} \rangle = \frac{\hbar^2}{2m} \left[\left(\frac{A}{c} \right)^2 (c-0) + \left(-\frac{A}{a-c} \right)^2 (a-c) \right] = +\frac{3\hbar^2}{2mc(a-c)} \quad (7)$$

which when combined with Eqn. (4) gives $\langle V_\delta \rangle + \langle \hat{T} \rangle = 0$, consistent with a zero-energy solution. To make use of the form in Eqn. (5), we note that $d\psi(x)/dx$ can be written in the form

$$\frac{d\psi(x)}{dx} = \frac{A}{c} + \left(-\frac{A}{a-c} - \frac{A}{c} \right) \Theta(x-c) \quad (8)$$

from which we find

$$\frac{d^2\psi(x)}{dx^2} = -\frac{Aa}{c(a-c)} \delta(x-c), \quad (9)$$

giving the same final result as in Eqn. (7). Either form can then be used to evaluate the expectation value of p^2 and the corresponding spread in momentum, namely

$$\langle p^2 \rangle = \frac{3\hbar^2}{c(a-c)} \quad \text{and} \quad \Delta p = \sqrt{\frac{3\hbar^2}{c(a-c)}}. \quad (10)$$

The momentum-space wavefunction for this system is also easily derived and is given by

$$\begin{aligned} \phi(p) &= \frac{1}{\sqrt{2\pi\hbar}} \int_0^a \psi(x) e^{-ipx/\hbar} dx \\ &= \sqrt{\frac{3}{2\pi\hbar a}} \left[\frac{\hbar^2}{p^2} \right] \left\{ \frac{1}{c} (e^{-ipc/\hbar} - 1) + \frac{1}{(a-c)} (e^{-ipc/\hbar} - e^{-ipa/\hbar}) \right\}. \end{aligned} \quad (11)$$

The corresponding probability density is then given by

$$\begin{aligned} |\phi(p)|^2 &= \frac{3\hbar^3}{\pi c(a-c)p^4} \left[\frac{1}{c} [1 - \cos(pc/\hbar)] + \frac{1}{(a-c)} [1 - \cos(p(a-c)/\hbar)] \right. \\ &\quad \left. - \frac{1}{a} [1 - \cos(pa/\hbar)] \right] \end{aligned} \quad (12)$$

which, despite the $1/p^4$ factor, is well-behaved at $p = 0$, giving $|\phi(0)|^2 = 3a/8\pi\hbar$, independent of c . The expectation value of p^2 can also be evaluated using this $|\phi(p)|^2$ to confirm the result in Eqn. (10). The momentum-space distribution, while giving the same $p = 0$ value for all values of c , does develop increasingly broad ‘wings’ as $c \rightarrow 0$ and $c \rightarrow a$ in order to achieve this result, as shown in Fig. 1.

More complex ZC waveforms can then be generated by appropriate combinations of multiple δ -functions. For example, the symmetric and antisymmetric (with respect to the center of the ISW) ZC wavefunctions

$$\psi^{(+)}(x) = \begin{cases} 0 & \text{for } x \leq 0 \text{ and } a \leq x \\ 3Ax/a & \text{for } 0 \leq x \leq a/3 \\ A & \text{for } a/3 \leq x \leq 2a/3 \\ 3A(1 - x/a) & \text{for } 2a/3 \leq x \leq a \end{cases} \quad (13)$$

(where $A = \sqrt{9/5a}$ for normalization) and

$$\psi^{(-)}(x) = \begin{cases} 0 & \text{for } x \leq 0 \text{ and } a \leq x \\ +3Bx/a & \text{for } 0 \leq x \leq a/3 \\ +3B(1 - 2x/a) & \text{for } a/3 \leq x \leq 2a/3 \\ -3B(1 - x/a) & \text{for } 2a/3 \leq x \leq a \end{cases} \quad (14)$$

(with $B = \sqrt{3/a}$) as shown in Fig. 2, can be obtained using a twin δ -function potential inside the ISW of the form

$$V_2(x) = g [\delta(x - a/3) + \delta(x - 2a/3)] \quad (15)$$

with critical strengths given by $g_0^{(+)} = -3\hbar^2/2ma$ and $g_0^{(-)} = -9\hbar^2/2ma$ respectively for these ZC solutions. These values are consistent with a stronger (more attractive) potential being required for the antisymmetric case to cancel the larger kinetic energy, as seen in Fig. 2. Generalizations to any arbitrary ‘straight-line’ waveform connecting very general sets of points of the form $(x, \psi(x))$ given by $(0, 0), (c_1, A_1), \dots, (c_N, A_N), (a, 0)$ are all possible with appropriately chosen combinations of additional δ -function potentials, provided one allows for the possibility of repulsive ($g > 0$) strengths. The kinetic/potential energy cancellation, yielding $\langle E \rangle = 0$, seen in the single δ -function potential case is easily generalizable to all such combinations.

An obvious limiting case for comparison is one in which any desired smooth waveform, $\psi_0(x)$, satisfying the ISW boundary conditions at $x = 0$ and $x = a$, can be approximated by a large number of short straight-line segments. These segments, defined by the points $(x_i, \psi_0(x_i))$, $i = 0, \dots, N$, where $N \rightarrow \infty$, would be matched appropriately using an increasingly large number of δ -functions. The resulting critical strengths are given by

$$g_i = \frac{\hbar^2}{2m} \left(\frac{\psi_0(x_i + \Delta x) - \psi_0(x_i)}{\Delta x} - \frac{\psi_0(x_i) - \psi_0(x_i - \Delta x)}{\Delta x} \right) / \psi_0(x_i) \approx \frac{\hbar^2}{2m} \frac{\psi_0''(x_i)}{\psi_0(x_i)} \quad (16)$$

so that the effective potential is given by

$$V_\delta(x) = \sum_i g_i \delta(x - x_i) \approx \frac{\hbar^2}{2m} \frac{\psi_0''(x)}{\psi_0(x)} \quad (17)$$

which reproduces the potential, $V_0(x)$, required to produce such an arbitrary $\psi_0(x)$ waveform with vanishing energy, namely

$$-\frac{\hbar^2}{2m} \frac{d^2 \psi_0(x)}{dx^2} + V_0(x) \psi_0(x) = 0. \quad (18)$$

Infinite square well plus δ -function systems have often been discussed in the context of wave packet revivals [5], [8], [9] where the time evolution of localized wave packets depends critically on the detailed energy spectrum, due to the $\exp(-iE_n t/\hbar)$ time dependence of the individual eigenstates used in the construction of any initial wave packet. Systems with identical or nearly identical energy spectra (isospectral) are then expected to exhibit

identical wave packet evolution and the simplest examples of such related systems are partner potentials related by supersymmetry [13]. It is useful to recall that in cases where a potential $V_{(-)}(x)$ is known to have a nodeless, ground state eigenstate given by $\psi_0(x)$, arranged to have zero energy ($E_0^{(-)} = 0$), then the supersymmetric partner potential given by

$$V_{(+)} = +V_{(-)}(x) - \frac{\hbar^2}{m} \frac{d}{dx} \left(\frac{\psi'_0(x)}{\psi_0(x)} \right) = -V_{(-)}(x) + \frac{\hbar^2}{m} \left(\frac{\psi'_0(x)}{\psi_0(x)} \right)^2. \quad (19)$$

The $V_{(\pm)}(x)$ partner potentials can also be written in terms of the superpotential, defined as

$$W(x) = -\frac{\hbar}{\sqrt{2m}} \left(\frac{\psi'_0(x)}{\psi_0(x)} \right), \quad (20)$$

in the form

$$V_{(\pm)}(x) = W^2(x) \pm \frac{\hbar}{\sqrt{2m}} W'(x). \quad (21)$$

The $V_{(\pm)}(x)$ potentials can be shown to have the identical energy spectra, except for the $E_0^{(-)} = 0$ state of $V_{(-)}(x)$ which is absent in the spectrum of $V_{(+)}(x)$; the energy eigenfunctions of $V_{(\pm)}(x)$, the $\psi_n^{(\pm)}(x)$, are then related to each other by generalized raising and lowering operators. (We are not considering periodic potentials in which it is possible [14] for both isospectral potentials to allow zero-energy ground state modes.)

Many of the ZC, and hence zero-energy, solutions discussed above are then immediate candidates for use in generating quite distinct partner potentials to the ISW plus δ -function potential systems considered here and elsewhere, which then have the same energy spectrum and presumably the same pattern of revivals, trapping, and bouncing discussed previously [7]. For example, the symmetric, ZC solution of the twin-delta function potential in Eqn. (13), with the critical value of $g_0^{(+)} = -3\hbar^2/2ma$, is the nodeless, zero-energy ground state of the potential $V_2(x)$ in Eqn. (15) and so can be used as the $\psi_0(x)$ in the construction of a supersymmetric partner potential. The corresponding $V_{(+)}(x)$ is given by

$$V_{(+)}(x) = +\frac{3\hbar^2}{2ma^2} [\delta(x - a/3) + \delta(x - 2a/3)] + \mathcal{V}(x) \quad (22)$$

where

$$\mathcal{V}(x) = \frac{\hbar^2}{2m} \begin{cases} 1/x^2 & \text{for } 0 < x < a/3 \\ 0 & \text{for } a/3 < x < 2a/3 \\ 1/(a-x)^2 & \text{for } 2a/3 < x < a \end{cases} \quad (23)$$

and this combination of two repulsive δ -function terms and the distinctive $\mathcal{V}(x)$ form will have the same overall spectrum as $V_2(x)$, except for the missing zero-energy ground state.

Finally, similar ZC solutions can be found for the infinite well modified with periodic boundary conditions (as considered in Refs. [7] and [8]) or in the related case of the quantum mechanical rotor defined by the Hamiltonian $\hat{H} = \hat{L}^2/2I$ with $\hat{L} \equiv (\hbar/i)(d/d\theta)$ where the eigensolutions are given by

$$\Theta_m(\theta) = \begin{cases} 1/\sqrt{2\pi} & \text{for } m = 0 \\ \cos(m\theta)/\sqrt{\pi} & \text{for } m > 0 \\ \sin(m\theta)/\sqrt{\pi} & \text{for } m > 0 \end{cases} \quad (24)$$

which are singly degenerate for $m = 0$ and doubly degenerate for $m \geq 1$. In these cases, ZC solutions are already present in the unperturbed system, corresponding in the case of the rotor to the $m = 0$ angular wavefunction given by $\Theta_0(\theta) = 1/\sqrt{2\pi}$. The addition of a single δ -function potential cannot provide a solution which is continuous in the periodic boundary case, but cases corresponding to the addition of two or more δ -functions can give a wide variety of ZC solutions. An example of one class of such solutions with two δ -functions is that in Eqn. (14), or any such case in which two equal strength δ -functions are placed symmetrically around the center of the standard infinite well, since this type of solution wraps around appropriately under periodic boundary conditions.

III. OTHER EXAMPLES

Piecewise-constant potentials also provide examples of local ZC wavefunctions for special choices of the well parameters. For example, the asymmetric infinite square well, defined by

$$V(x) = \begin{cases} +\infty & \text{for } x < -a \\ 0 & \text{for } -a < x < 0 \quad (\text{Region I}) \\ +V_0 & \text{for } 0 < x < +b \quad (\text{Region II}) \\ +\infty & \text{for } +b < x \end{cases} \quad (25)$$

can support energy eigenfunctions corresponding to both $E > V_0$ and $V_0 > E > 0$ and the conditions on the energy eigenvalues for each case are easily derived to be

$$\begin{aligned} q \tan(ka) + k \tan(qb) &= 0 & \text{with } k &\equiv \sqrt{\frac{2mE}{\hbar^2}} & \text{and } q &\equiv \sqrt{\frac{2m(E - V_0)}{\hbar^2}} \\ \kappa \tan(ka) + k \tanh(\kappa b) &= 0 & \text{where } \kappa &\equiv \sqrt{\frac{2m(V_0 - E)}{\hbar^2}} \end{aligned} \quad (26)$$

for $E > V_0$ and $V_0 > E > 0$ solutions respectively. Turning the problem around, either form can then be used to find the condition on the well parameters necessary to have a zero curvature solution with $E = V_0$, namely, by taking the $q \rightarrow 0$ or $\kappa \rightarrow 0$ limits where one finds

$$\tan(\chi a) = -\chi b \quad \text{with} \quad \chi \equiv \sqrt{\frac{2mV_0}{\hbar^2}}. \quad (27)$$

In this case, the solutions are given by

$$\psi(x) = \begin{cases} A \sin[\chi(x + a)] & \text{for } -a < x < 0 \\ C(1 - x/b) & \text{for } 0 < x < +b \end{cases} \quad (28)$$

with ZC solutions in the right side of the well. For large values of χ , the condition on the well parameters in Eqn. (27) reduces to $\chi \approx (2n + 1)\pi/2a$ (an integral number of half wavelengths plus an extra quarter wavelength in the left side of the well). The probabilities that the particle is found in the left (I) and right (II) sides of the well approach

$$P_I = \frac{3a}{(3a + 2b)} \quad \text{and} \quad P_{II} = \frac{2b}{(3a + 2b)}. \quad (29)$$

Supersymmetric partner potentials for the asymmetric infinite well are also easy to construct in the case where the ZC solution is the nodeless ground state of the system.

Many similar examples are possible in such piecewise-constant potentials, including regions where the ZC wave function is actually flat (in the case of a symmetric infinite well with an enclosed symmetric barrier of height V_0) or as a threshold solution of the finite square well, consistent with discussions of Ref. [12] in which the poles of the S -matrix for the finite square well are analyzed for bound, threshold, and scattering states. The analyses of all such cases is straightforward and will be discussed in a more pedagogical context elsewhere [15].

Acknowledgments We would like to thank Laura P. Gilbert for her helpful feedback in the preparation of this manuscript. The work of MB was supported in part by a Research Corporation Cottrell College Science Award (CC5470).

-
- [1] V. V. Nesvizhevsky *et al.*, Nature **415**, 297-299 (2002); Phys. Rev. D **67**, 102002 (2003); *ibid*, **68**, 108702 (2003).
 - [2] D. M. Meekhof, C. Monroe, B. E. King, W. M. Itano, and D. J. Wineland, Phys. Rev. Lett. **76**, 1796 (1996).
 - [3] A. Bonvalet, J. Nagle, V. Berger, A. Migus, J. -L. Martin, and M. Joffre, Phys. Rev. Lett. **76**, 4392 (1996).
 - [4] V. Venugopalan and G. S. Agarwal, Phys. Rev. A, **59**, 1413 (1999); D. L. Aronstein, and C. R. Stroud, Jr., Phys. Rev. A **62**, 022102 (2000); Am. J. Phys. **68**, 943 (2000).
 - [5] G. A. Vugalter, A. K. Das, and V. A. Sorokin, Phys. Rev. A **66**, 012104 (2002).
 - [6] A. C. de la Torre, H. O. Martín, and D. Goyeneche, Phys. Rev. E **68**, 031103 (2003).
 - [7] A. G. M. Schmidt, B. K. Cheng, and M. G. E. da Luz, Phys. Rev. A **66**, 062712 (2002).
 - [8] A. G. M. Schmidt and M. G. E. da Luz, Phys. Rev. A **69**, 052708 (2004).
 - [9] C. U. Segre and J. D. Sullivan, Am. J. Phys. **44**, 729 (1976). This paper was one of the first to describe wave packet revivals, as well as to consider the effect of δ -function barriers on such behavior in the context of the infinite square well.
 - [10] U. Kuhl and H.-J. Stöckman, Physics E (Amsterdam) **9**, 384 (2001); U. Kuhl, F. M. Izrailev, A. A. Krokhnin, and H.-J. Stöckman, Appl. Phys. Lett. **77**, 633 (2000); U. Kuhl and H.-J. Stöckman, Phys. Rev. Lett. **80**, 3232 (1998).
 - [11] One can, of course, solve for the quantized energy eigenvalues and normalized wavefunctions for general $E > 0$ and $E < 0$ solutions with the additional δ -function interaction, and both yield the results here in the limit that $E \rightarrow 0$.
 - [12] D. W. L. Sprung, H. Wu, and J. Martorell, Am. J. Phys. **64**, 136 (1996).
 - [13] For a review of supersymmetric quantum mechanics, see F. Cooper, A. Khare, and U. Sukhatme, Phys. Rep. **251**, 267 (1995); *Supersymmetry in quantum mechanics*, World Scientific Press, Singapore (2002).
 - [14] G. Dunne and J. Feinberg, Phys. Rev. D **57**, 1271 (1998).
 - [15] L. P. Gilbert, M. Belloni, M. A. Doncheski, and R. W. Robinett, to be submitted for publication.




FIG. 1: Plots of $\psi(x)$ versus x for two zero-curvature wavefunctions in the infinite square well plus δ -function potential system (left) with the additional $V_\delta(x; c)$ potential located at $c = a/2$ (dashed lines) and $c = 0.01a$ (solid lines). The corresponding momentum-space probability densities, $|\phi(p)|^2$ versus p (in units of $\hbar/a$), are shown on the right.




FIG. 2: Plots of $\psi(x)$ versus x for the symmetric (solid) and antisymmetric (dashed) zero-curvature states in Eqs. (13) and (14) respectively.